# Journal **CHEP**

09-09-05

**Eric F Post** 

Division of Otorhinolaryngology Faculty of Health Sciences Tygerberg Campus, University of Stellenbosch

# Supracricoid Partial Laryngectomy (larynx Ca)

- Italy
- Retrospective
- Aim: 1. Evaluate oncological outcome2. Determine prognostic factors
- 253 pt = 180 CHP glottic Ca 138
  SG Ca 42
  - 73 CHEP glottic

Table 1. Pathological TNM Classification for CHP and CHEP

Pathological TNM Classification	CHP		CHEP	
	Glottic (n = 138)	Supraglottic (n = 42)	Glottic (n = 73)	Total, % (N = 253)
T1 N0	6	2	19	10.7
T2 N0	68	22	48	54.5
T2 N1	4	2	0	2.4
T2 N2	2	1	0	1.2
T3 N0	45	7	5	22.5
T3 N1	2	3	1	2.4
T3 N2	1	0	0	0.4
T4 N0	7	3	0	4.0
T4 N1	1	0	0	0.4
T4 N2	2	2	0	1.6

Abbreviations: CHEP, cricohyoidoepiglottopexy; CHP, cricohyoidopexy

### CHEP


- Cricohyoidepiglottopexy
- Preserve swallow + speech
  - Spare part of epiglottis, arytenoids with innervation
- Sleeve-like resection:
  - Thyroid cartilages + base of epiglottis + paraglottic space removed
  - From hyoid cricoid


#### CHEP


- Piquet et al, France, 1974
- T2, T3 glottic Ca
- 75 % 5 yr survival
- 5% recurrence
- All N deglutition (2 occ. Aspiration)
- All satisfactory speech
  - Arytenoid mucosa vibrate against epiglottis


## CHEP

- Indications
  - Ant 2/3 VC
  - Entire VC with ↓ mobility
  - Early ventricular Ca
  - Early ant. Commisure Ca
  - Both VC
- Contraindications
  - Immobile arytenoid (tumor fixing cartilage)
  - Subglottic > 7 mm









#### Results

- (+) margins
  - Dysplasia = invasive Ca
    - Timing differs
  - 0.5 cm or less from margin
  - Only predictor of recurrence
- 5 yr survival
  - **86,1 %**
  - Recurrence 45,5%

TL +/- RoRx

#### Results

- SCPL = valid Rx for selective group of pt
  - CHEP alt to partial L (vertical / horisontal)
  - CHP alt to TL

- Prognostic factors for survival
  - 1<sup>st</sup> = Recurrence
  - Also T, N, staging, (+) margins